

Community-Acquired *Listeria monocytogenes* Meningitis in Adults

Matthijs C. Brouwer,¹ Diederik van de Beek,¹ Sebastiaan G. B. Heckenberg,¹ Lodewijk Spanjaard,² and Jan de Gans¹

Departments of ¹Neurology and ²Medical Microbiology, Center of Infection and Immunity Amsterdam, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Background. *Listeria monocytogenes* is the third most common cause of bacterial meningitis.

Methods. We prospectively evaluated 30 episodes of community-acquired *L. monocytogenes* meningitis, confirmed by culture of cerebrospinal fluid specimens, in a nationwide study in The Netherlands. Outcome was graded using the Glasgow outcome score; an unfavorable outcome was defined as a score of 1–4.

Results. We found 30 episodes of *L. monocytogenes* meningitis. All patients were immunocompromised or >50 years old. In 19 (63%) of 30 patients, symptoms were present for >24 h; in 8 patients (27%), symptoms were present for ≥4 days. The classic triad of fever, neck stiffness, and change in mental status was present in 13 (43%) of 30 patients. An individual cerebrospinal fluid indicator of bacterial meningitis was present in 23 (77%) of 30 cases. Gram staining of cerebrospinal fluid samples revealed the causative organism in 7 (28%) of 25 cases. The initial antimicrobial therapy was amoxicillin based for 21 (70%) of 30 patients. The coverage of initial antimicrobial therapy was microbiologically inadequate for 9 (30%) of the patients. The mortality rate was 17% (5 of 30), and 8 (27%) of 30 patients experienced an unfavorable outcome. Inadequate initial antimicrobial therapy was not related to outcome.

Conclusions. In contrast with previous reports, we found that patients with meningitis due to *L. monocytogenes* do not present with atypical clinical features; however, typical cerebrospinal fluid findings predictive for bacterial meningitis might be absent. A high proportion of patients received initial antimicrobial therapy that did not cover *L. monocytogenes*.

Bacterial meningitis is a serious and life-threatening disease. The estimated incidence is 4–6 cases per 100,000 adults per year in developed countries and is up to 10 times higher in less developed countries [1, 2]. *Streptococcus pneumoniae* and *Neisseria meningitidis* are the leading causes of bacterial meningitis in adults and cause 85% of all cases [2, 3]. *Listeria monocytogenes* is the third most frequent cause of bacterial meningitis [2–5]. This gram-positive bacillus is principally spread by contaminated food, which was discovered after outbreaks of listeriosis in the 1980s [1, 6–9]. In the literature, meningitis due to *L. monocytogenes* is described as a disease in immunocompromised patients and el-

derly individuals, with high case-fatality rates (24%–62%) [1, 5, 10–17]; however, all previous data on *Listeria* meningitis in adults were retrospective. We performed the first prospective nationwide cohort study in adults with community-acquired bacterial meningitis [2]. In this report, we describe the clinical features, complications, treatment, and outcome in adults with bacterial meningitis due to *L. monocytogenes* in this prospective case series.

METHODS

In the Dutch Meningitis Cohort Study, a nationwide observational cohort study in The Netherlands, 696 episodes of community-acquired acute bacterial meningitis were assessed prospectively. All causative organisms were identified by CSF culture, which yielded *S. pneumoniae* in 352 episodes (51%), *N. meningitidis* in 257 (37%), *L. monocytogenes* in 30 (4%), and other bacteria in 57 (8%). Inclusion and exclusion criteria have been described elsewhere [2]. In summary, eligible patients were >16 years old, had bacterial meningitis

Received 8 May 2006; accepted 14 June 2006; electronically published 10 October 2006.

Reprints or correspondence: Dr. Matthijs C. Brouwer, Academic Medical Center, University of Amsterdam, Dept. of Neurology H2, P.O. Box 22660, 1100 DD Amsterdam, The Netherlands (m.c.brouwer@amc.uva.nl).

Clinical Infectious Diseases 2006;43:1233–8

© 2006 by the Infectious Diseases Society of America. All rights reserved.
1058-4838/2006/4310-0001\$15.00

Table 1. Clinical and laboratory characteristics at admission to the hospital for 30 adults with community-acquired *Listeria monocytogenes* meningitis.

Variable	Patients (n = 30)
Age, mean years \pm SD	65 \pm 18
Male sex	15/30 (50)
Predisposing factors	
Immunocompromise	20/30 (67)
Pneumonia	1/30 (3)
Otitis or sinusitis	1/30 (3)
Pretreated with antimicrobials	5/30 (17)
Duration of symptoms <24 h	11/30 (37)
Seizures	2/30 (7)
Symptoms at presentation	
Headache	22/25 (88)
Nausea	20/24 (83)
Neck stiffness	22/30 (73)
Temperature \geq 38°C	27/30 (90)
Score on Glasgow coma scale at presentation	
Median score \pm SD	12 \pm 3
<14 (indicating change in mental status)	21/30 (70)
\leq 8 (indicating coma)	3/30 (10)
Triad of fever, neck stiffness, and change in mental status	13/30 (43)
Focal neurological deficits	
Any	11/30 (37)
Aphasia	7/25 (28)
Hemiparesis	2/30 (7)
Cranial nerve palsies	2/30 (7)
Ataxia	2/30 (7)
Laboratory findings ^a	
Indexes of CSF inflammation	
Opening pressure, median mm of water (range)	275 (150–400)
WBC count	
Median cells (range)	620 (24–16003)
<100 cells/mL	4/28 (13)
100–999 cells/mL	13/28 (46)
>999 cells/mL	11/28 (39)
Protein level, median g/L (range)	2.52 (1.1–19.3)
CSF/blood glucose ratio, median value (range)	0.30 (0.03–0.86)
CSF Gram stain results	
Negative	16/25 (60)
Gram-positive rod	7/25 (28)
Other ^b	2/25 (4)
Blood culture findings	
<i>L. monocytogenes</i>	12/26 (46)
Other ^c	2/26 (8)
Erythrocyte sedimentation rate, median mm per h (range)	39 (10–242)
C-reactive protein level, median value (range)	117 (4–467)
Thrombocyte count, median platelets/mL (range)	186 (4–653)
Hyponatremia	22/30 (73)

NOTE. Data are proportion (%) of patients, unless otherwise indicated.

^a CSF pressure was measured in 8 patients, CSF WBC count in 28, CSF protein concentration in 29, CSF/blood glucose ratio in 28, erythrocyte sedimentation rate in 25, C-reactive protein level in 22, and the thrombocyte count in 30.

^b Gram-negative bacteria in 2 patients.

^c *Streptococcus* and *Staphylococcus* species.

Table 2. Demographic and clinical characteristics of immunocompetent patients with *Listeria monocytogenes* meningitis.

Patient	Age, years	Sex	Duration of symptoms <24 h	Classic triad of symptoms	Outcome
1	87	F	Yes	Absent	Death
2	65	M	No	Absent	Good recovery
3	90	F	Yes	Present	Good recovery
4	72	F	Yes	Absent	Good recovery
5	79	F	Yes	Present	Good recovery
6	58	M	No	Present	Good recovery
7	77	F	Yes	Absent	Good recovery
8	56	M	Yes	Present	Death
9	69	F	Yes	Absent	Good recovery
10	73	F	Yes	Absent	Good recovery

confirmed by culture of CSF, and were listed in the database of The Netherlands Reference Laboratory for Bacterial Meningitis from October 1998 to April 2002. This laboratory provided daily updates of the names of hospitals where patients with bacterial meningitis had been admitted 2–6 days previously. The start of the cohort study was announced in the journal of the Dutch Neurological Society, with periodic reminders. Before the study started, all neurologists received information by mail about the study, including a case record form. For each patient, the treating physician was contacted, and informed consent was obtained from all participating patients or their legally authorized representatives. The Dutch Meningitis Cohort Study was approved by the ethics committee of the Academic Medical Center, University of Amsterdam (Amsterdam, The Netherlands).

Data registration was started at the time of inclusion. The study was undertaken in accordance with Dutch privacy legislation, and information was obtained via a case-record form. Patients receiving immunosuppressive drugs and patients with diabetes mellitus, alcoholism, asplenia, liver cirrhosis, end-stage renal disease, or HIV infection were judged to be immunocompromised. Persisting fever was defined as a body temperature of $\geq 38^{\circ}\text{C}$ for >10 days after the start of appropriate antimicrobial treatment. Recurrent fever was defined as fever reoccurring after at least 1 afebrile day. Hyponatremia was defined as a plasma sodium concentration <135 mmol/L. Patients underwent a neurological examination at discharge, and outcome was graded with the Glasgow outcome scale. This measurement scale is well validated, with scores varying from 1 (death) to 5 (good recovery). A favorable outcome was defined as a score of 5, and an unfavorable outcome was defined as a score of 1–4. Focal neurological abnormalities were divided into focal cerebral deficits (aphasia, monoparesis, or hemiparesis) and cranial nerve palsies. Causes of death were independently classified by 2 investigators (M.C.B. and D.v.d.B.) in 2 categories on the basis of criteria described in previous studies [18,

19]. These 2 categories were: (1) systemic causes, including septic shock, respiratory failure, multiorgan dysfunction, and cardiac ischemia; and (2) neurological causes, including brain herniation, cerebrovascular complications, intractable seizures, and withdrawal of care because of poor neurological prognosis. There were no differences in classification between clinicians.

The Mann-Whitney *U* test was used to identify differences between groups with respect to continuous variables, and dichotomous variables were compared with use of the χ^2 test. All statistical tests were 2-tailed, and a *P* value of <.05 was regarded as significant. Analyses were undertaken with SPSS software, version 12.0.1 (SPSS).

RESULTS

In a 3.5-year period, 30 episodes of community-acquired *L. monocytogenes* meningitis were identified in 30 patients (table 1). The calculated annual incidence was 0.07 cases per 100,000 adults. There was no seasonal pattern of occurrence of *L. monocytogenes* meningitis, although 5 (50%) of 10 episodes in immunocompetent patients occurred during the summer. Predisposing conditions were present in 21 (70%) of 30 patients; 20 (67%) of 30 patients were considered to be immunocompromised, and 1 patient had sinusitis (3%). All patients who were characterized as immunocompetent were >50 years old (table 2). A previous episode of meningitis had occurred in 1 patient (3%). In 19 (63%) of 30 patients, symptoms were present for >24 h; in 8 (27%), symptoms were present for ≥ 4 days.

Classic symptoms and signs were present in most of the patients. The classic triad of neck stiffness, fever, and altered mental status was found in 13 (43%) of 30 patients. At least 2 of 4 symptoms and signs (classic triad plus headache) were present in 29 (97%) of the patients. One patient had a rash that was characterized as ecchymosis.

CT of the brain was performed at admission for 23 (77%)

of 30 patients; the results were normal for 18 patients. Abnormalities were recorded for 5 patients (27%). Cerebral edema was present in 2 patients (9%), evidence of recent infarction was present in 1 patient (5%), lesions possibly indicating an infarction at an earlier date were present in 1 patient (5%), and atrophy was present in 2 patients (9%). Cranial CT was performed before lumbar puncture for 16 (70%) of the 23 patients; therapy was started before CT for only 5 (31%) of the patients. MRI of the brain was performed for 1 patient and revealed no abnormalities.

Lumbar puncture was performed for all patients. Opening pressure was recorded with a water-manometer for 8 (27%) of 30 patients; median pressure was 275 mm of water (range, 150–400 mm of water), and 5 patients had very high pressure (>250 mm of water). At least 1 individual CSF finding indicative of bacterial meningitis (glucose concentration, <1.9 mmol/L; ratio of CSF glucose to blood glucose, <0.23; protein concentration, >2.20 g/L; WBC count, >2000 cells/mL; or CSF neutrophil count, 1180 cells/mL [20]) was found for 23 (77%) of 30 patients. There was no relation between low CSF leukocyte count and age or immunocompetence. Gram staining of CSF samples was performed for 25 (83%) of 30 patients and revealed no microorganisms in 16 (64%), gram-positive rods in 7 (28%), and gram-negative bacteria in 2 (8%).

Initial antimicrobial treatment consisted of penicillin or amoxicillin for 9 (30%) of 30 patients, a third-generation cephalosporin for 5 (17%), and a combination of amoxicillin and a third-generation cephalosporin for 8 (27%); other regimens were used for 8 patients (27%). In 21 episodes (70%), the initial therapy included penicillin or amoxicillin; a combination with aminoglycosids was given in 7 (23%) of the episodes. Nine (30%) of 30 patients received inappropriate initial antimicrobial therapy. Among these patients, the median time from admission to the hospital to receipt of adequate therapy was 3 days (range, 1–5 days). Adjunctive steroids were given in 5 episodes (17%); however, 3 patients had already used prednisone before admission to the hospital, and 1 patient received dexamethasone as treatment for an exacerbation of chronic obstructive pulmonary disease. In 1 additional patient, dexamethasone therapy was started after the patient experienced clinical deterioration.

Complications developed in a high proportion of patients (table 3). Persisting fever was present in 11 (37%) of 30 patients. Subsequent lumbar punctures were performed for 3 patients, which all showed a decrease in CSF pleocytosis; CSF cultures had negative results for all patients. The cause of persisting fever was unknown in most patients (63%); other causes were pneumonia in 2 (18%), drug-related fever in 1 (9%), and sepsis in 1 (9%). Mean time to defervescence was 4.9 days (range, 1–18 days). Recurrent fever occurred in 11 episodes (37%); causes were unknown in 6 (54%), pneumonia in 3 (27%), and drug-

Table 3. Complications during hospitalization and outcome for patients with *Listeria monocytogenes* meningitis.

Variable	Proportion of patients (%)
Complication	
Seizures	6/30 (20)
Cardiorespiratory failure	10/30 (33)
Receipt of mechanical ventilation	7/30 (23)
Sepsis	5/30 (17)
Hyponatremia	25/30 (83)
Recurrent fever	11/28 (39)
Persisting fever	11/30 (37)
Impaired consciousness	12/30 (40)
Other complications ^a	7/30 (23)
Outcome	
Score on the Glasgow outcome scale	
1 (death)	5/30 (17)
2 (vegetative state)	0/30 (0)
3 (severe disability)	1/30 (3)
4 (moderate disability)	2/30 (7)
5 (mild or no disability)	22/30 (73)
Sequelae at discharge	
Hemiparesis	2/25 (8)
Cranial nerve palsy ^b	2/25 (8)

^a Other complications included pneumonia in 3 (10%) of patients, drug-related fever in 2 (7%), alcohol abstinence delirium in 1 (3%), and joint pain in 1 (3%).

^b Trochlear and facial nerve palsy each occurred in 1 patient.

related fever in 2 (18%). During hospitalization, cranial CT scanning was repeated for 5 patients; 1 of these CT scans had abnormal findings, showing a subarachnoid hemorrhage.

Hyponatremia was found at admission to the hospital for 22 (73%) of 30 patients; 8 patients had severe hyponatremia (plasma sodium level, <130 mmol/L), and 14 patients had mild hyponatremia (plasma sodium level, <135 mmol/L). An additional 3 patients developed hyponatremia during hospitalization. Five (20%) of 25 patients were treated for hyponatremia; in 1 patient, this treatment consisted of fluid restriction.

Five (17%) of 30 patients in this cohort died. All of these patients died within 3 days after admission to the hospital, and all patients who died were >55 years old. Two of these 5 patients died of systemic complications, and 3 patients died of neurological complications. One patient died following a subarachnoid hemorrhage 1 day after admission to the hospital. Neurological examination was performed at discharge from the hospital for all 25 surviving patients and revealed cranial nerve palsies in 2 (8%) and focal cerebral deficits in 2 (8%). No patients had hearing impairment. In a univariate analysis, predictors for an unfavorable outcome were a low score on the Glasgow coma scale at admission to the hospital ($P = .015$) and a low platelet count ($P = .024$). Of the 9 patients who received microbiologically inadequate initial antimicrobial ther-

apy, 2 (22%) of died; the other 7 patients had favorable outcomes.

DISCUSSION

Our findings show that meningitis due to *L. monocytogenes* is a disease that occurs among immunocompromised patients and elderly individuals. This corresponds with previous reports, which describe *L. monocytogenes* infection in patients with impaired cellular immunity (e.g., attributable to immunosuppressive therapy, immunosenescence, diabetes, or malignancies) [23, 24]. *L. monocytogenes* meningitis in young, previously healthy adults has been reported only in anecdotal observations [11, 14]. In a large literature review including all case series and case reports, this group constitutes only 6% of patients with *L. monocytogenes* CNS infection [14]. In our series, all patients were either immunocompromised or >50 years old.

Symptoms and signs of patients presenting with *L. monocytogenes* meningitis were not different from those found in the general population of patients with community-acquired bacterial meningitis. The prevalence of the classic triad of fever, neck stiffness, and altered mental status was 43%, and almost all patients presented with at least 2 of the 4 classic symptoms of headache, fever, neck stiffness, and altered mental status. Similar prevalences have been reported in the general population of patients with bacterial meningitis [2, 21]. This is in contrast with previous reports, which stressed the importance of the atypical presentation of patients with *L. monocytogenes* meningitis [14, 22]. This difference may well be explained by the retrospective design of these previous studies. This is the first prospective study of *L. monocytogenes* meningitis to date. Concordant with previous reports was the subacute development of disease in most cases [14, 22]. In a large proportion of patients, symptoms were present for ≥ 4 days (27%).

A high proportion of adults with *L. monocytogenes* meningitis had atypical CSF findings; 23% of the patients had no individual CSF findings indicative of bacterial meningitis. In addition, Gram staining had a low yield. The causative microorganism was revealed by Gram staining in only 24% of patients. The atypical CSF findings and the low yield of Gram staining have been described previously in *L. monocytogenes* meningitis [3, 14].

Guidelines for suspected bacterial meningitis recommend amoxicillin-based empirical antimicrobial therapy for patients aged >50 years old or with risk factors to cover *L. monocytogenes*, because this bacterium is resistant to cephalosporins [25, 26]. Most patients in our case series received empirical therapy, including amoxicillin (70%). However, a considerable proportion of patients received third-generation cephalosporin monotherapy (27%). The median delay before these patients received adequate antimicrobial therapy was 3 days. This delay, however, was not associated with an unfavorable outcome. Although

speculative, the absence of a relation between delay of adequate therapy and outcome might be explained by the limited number of patients included in the study. Another explanation could be the subacute development of disease in patients with *L. monocytogenes* meningitis. If meningitis due to *L. monocytogenes* is proven, either by Gram stain or CSF culture, we advise that patients be treated with amoxicillin, administered at a dosage of 2 g every 4 h for at least 21 days [22, 25, 27–29].

A clinical trial showed a beneficial effect of early dexamethasone treatment in adults with bacterial meningitis [30]. A recent meta-analysis confirmed these results [31]. Although 5 patients in our case series were eventually treated with steroids, none received adjunctive dexamethasone according to current guidelines [25]. Immunocompromised patients were excluded from the studies of adjunctive dexamethasone [30, 31]; therefore, these patients should not receive dexamethasone. Dexamethasone treatment is indicated for most patients with suspected bacterial meningitis; however, it remains unclear whether dexamethasone has beneficial effects in *L. monocytogenes* meningitis.

Our study has several important limitations. First, only patients who had a positive CSF culture result were included. Negative CSF culture results occur in 11%–30% of patients with bacterial meningitis, and this rate may be even higher among patients with *L. monocytogenes* meningitis, although exact figures are unknown [2, 3, 13, 22]. However, the clinical presentation of patients with positive and negative CSF culture results was closely similar in several studies [3, 5, 24]. Furthermore, patients with space-occupying lesions visible on CT may not undergo lumbar puncture [26]. Brain abscesses are found in 5% of *L. monocytogenes* infections of the CNS [14]. In patients with coagulation disorders or severe septic shock, lumbar puncture might be postponed, which can result in CSF cultures with negative results, as well [25]. Therefore, these patient groups were probably only partly represented in our study.

We describe, to our knowledge, the first patient with a subarachnoid hemorrhage as a complication in *L. monocytogenes* meningitis. Subarachnoid hemorrhage is known to be a rare complication of tuberculous meningitis, actinomycotic meningitis, and bacterial meningitis due to *S. pneumoniae*, but it has never been described in *L. monocytogenes* meningitis [32–34].

Another interesting observation in our study was the high rate of patients with hyponatremia (83%). Although high rates of hyponatremia are known from case series of tuberculous and group A streptococci meningitis, this has not been described in *L. monocytogenes* meningitis [35, 36]. Probable causes of hyponatremia are the syndrome of inappropriate antidiuretic hormone secretion and the cerebral salt wasting syndrome [37, 38]. The appropriate treatment for syndrome of inappropriate

antidiuretic hormone secretion—fluid restriction—is opposite from the usual treatment of hyponatremia caused by cerebral salt wasting syndrome. Differentiation between the syndromes, however, is difficult. In children with bacterial meningitis and hyponatremia, fluid restriction does not improve either brain edema or outcome [39]. Current guidelines recommend that treatment of adults with bacterial meningitis should aim for a normovolemic state [25].

In conclusion, our study showed that *L. monocytogenes* meningitis is found in immunosuppressed patients and elderly individuals. Patients presented with signs and symptoms that were similar to those of the general population with bacterial meningitis, albeit with a longer prodromal phase. Despite frequent complications during hospitalization, the associated mortality rate was relatively low.

Acknowledgments

We are indebted to many physicians in The Netherlands for their cooperation.

Financial support. Baxter BV (research grant to J.d.G.), the Meningitis Research Foundation (research grant to D.v.d.B.), and The Netherlands Organization for Health Research and Development (ZonMw) (NWO-Veni grant 2006 [916.76.023] and NWO-Rubicon grant 2006 [019.2006.1.310.0001] to D.v.d.B.). The Dutch Meningitis Cohort Study was supported in part by a research grant from Roche Pharmaceuticals.

Potential conflicts of interest. All authors: no conflicts.

References

- Schlech WF, Ward JI, Band JD, et al. Bacterial meningitis in the United States, 1978 through 1981: the national bacterial meningitis surveillance study. *JAMA* **1985**;253:1749–54.
- van de Beek D, de Gans J, Spanjaard L, et al. Clinical features and prognostic factors in adults with bacterial meningitis. *N Engl J Med* **2004**;351:1849–59.
- Durand ML, Calderwood SB, Weber DJ, et al. Acute bacterial meningitis in adults. *N Engl J Med* **1993**;328:21–8.
- Wenger JD, Hightower AW, Facklam RR, et al. Bacterial meningitis in the United States, 1986: report of a multistate surveillance study. *J Infect Dis* **1990**;162:1316–23.
- Sigurdardottir B, Bjornsson OM, Jonsdottir KE, et al. Acute bacterial meningitis in adults: a 20-year overview. *Arch Intern Med* **1997**;157:425–30.
- Bula CJ, Bille J, Glauser MP. An epidemic of food-borne listeriosis in western Switzerland: description of 57 cases involving adults. *Clin Infect Dis* **1995**;20:66–72.
- Schuchat A, Robinson K, Wenger MD, et al. Bacterial meningitis in the united states in 1995. *N Engl J Med* **1997**;337:970–6.
- Fleming DW, Cochi SL, MacDonald KL, et al. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. *N Engl J Med* **1985**;312:404–7.
- Linnan MJ, Mascola L, Lou XD, et al. Epidemic listeriosis associated with Mexican-style cheese. *N Engl J Med* **1988**;319:823–8.
- Pigrau C, Almirante B, Pahissa A, et al. Clinical presentation and outcome in cases of listeriosis. *Clin Infect Dis* **1993**;17:143–4.
- Cherubin CE, Marr JS, Sierra MF, et al. Listeria and gram-negative bacillary meningitis in New York City, 1972–1979: frequent causes of meningitis in adults. *Am J Med* **1981**;71:199–209.
- Skogberg K, Syrjanen J, Jahnkola M, et al. Clinical presentation and outcome of listeriosis in patients with and without immunosuppressive therapy. *Clin Infect Dis* **1992**;14:815–21.
- Gellin BG, Broome CV. Listeriosis. *JAMA* **1989**;261:1313–20.
- Mylonakis E, Hohmann EL, Calderwood SB. Central nervous system infection with *Listeria monocytogenes*: 33 years of experience at a general hospital and review of 776 episodes from the literature. *Medicine Baltimore* **1998**;77:313–36.
- Calder JAM. Listeria meningitis in adults. *Lancet* **1997**;350:307–8.
- Aouaj Y, Spanjaard L, van Leeuwen N, et al. *Listeria monocytogenes* meningitis: serotype distribution and patient characteristics in the Netherlands, 1976–95. *Epidemiol Infect* **2002**;128:405–9.
- Hussein AS, Shafran SD. Acute bacterial meningitis in adults: a 12-year review. *Medicine Baltimore* **2000**;79:360–8.
- McMillan DA, Lin CY, Aronin SI, et al. Community-acquired bacterial meningitis in adults: categorisation of causes and timing of death. *Clin Infect Dis* **2001**;33:969–75.
- van de Beek D, de Gans J. Dexamethasone and pneumococcal meningitis. *Ann Intern Med* **2004**;141:327.
- Spanos A, Harrell FE Jr, Durack DT. Differential diagnosis of acute bacterial meningitis: an analysis of the predictive value of initial observations. *JAMA* **1989**;262:2700–7.
- Attia J, Hatala R, Cook DJ, et al. The rational clinical examination: does this adult patient have acute meningitis? *JAMA* **1999**;282:175–81.
- Lorber B. Listeriosis. *Clin Infect Dis* **1997**;24:1–11.
- Lorber B. *Listeria monocytogenes*. In: Mandell GL, Bennet JL, Dolin R, eds. *Mandell, Douglas, and Bennet's principles and practice of infectious disease*. 5th ed. New York: Churchill Livingstone, **2000**;2208–15.
- Tunkel AR. Bacterial meningitis. Philadelphia, PA: Lippincott Williams & Wilkins, **2000**.
- van de Beek D, de Gans J, Tunkel AR, et al. Community-acquired bacterial meningitis in adults. *N Engl J Med* **2006**;354:44–53.
- van de Beek D, de Gans J, Spanjaard L, et al. Antibiotic guidelines and antibiotic use in adults: bacterial meningitis in the Netherlands. *J Antimicrob Chemother* **2002**;49:661–6.
- Southwick FS, Purich DL. Intracellular pathogenesis of listeriosis. *N Engl J Med* **1996**;334:770–6.
- Quagliarello VJ, Scheld WM. Treatment of bacterial meningitis. *N Engl J Med* **1997**;336:708–16.
- Hof H. Listeriosis: therapeutic options. *FEMS Immunol Med Microbiol* **2003**;35:203–5.
- de Gans J, van de Beek D. Dexamethasone in adults with bacterial meningitis. *N Engl J Med* **2002**;347:1549–56.
- van de Beek D, de Gans J, McIntyre P, et al. Steroids in adults with acute bacterial meningitis: a systematic review. *Lancet Infect Dis* **2004**;4:139–43.
- Kastenbauer S, Pfister HW. Pneumococcal meningitis in adults: spectrum of complications and prognostic factors in a series of 87 cases. *Brain* **2003**;126:1015–25.
- Yeh ST, Lee WJ, Lin HJ, et al. Nonaneurysmal subarachnoid hemorrhage secondary to tuberculous meningitis: report of two cases. *J Emerg Med* **2003**;25:265–70.
- Koda Y, Seto Y, Takeichi S, et al. Fatal subarachnoid hemorrhage complicating actinomycotic meningitis. *Forensic Sci Int* **2003**;134:169–71.
- Haas DW. Mycobacterial diseases. In: Mandell GL, Bennet JL, Dolin R, eds. *Mandell, Douglas, and Bennet's principles and practice of infectious disease*. 5th ed. New York: Churchill Livingstone, **2000**;2598–9.
- van de Beek D, de Gans J, Spanjaard L, et al. Group A streptococcal meningitis in adults: report of 41 cases and a review of the literature. *Clin Infect Dis* **2002**;34:e32–6.
- Harrigan MR. Cerebral salt wasting syndrome: a review. *Neurosurgery* **1996**;38:152–60.
- Coenraad MJ, Meinders AE, Taal JC, et al. Hyponatremia in intracranial disorders. *Neth J Med* **2001**;58:123–7.
- Oates-Whitehead RM, Maconochie I, Baumer H, et al. Fluid therapy for acute bacterial meningitis. *Cochrane Database Syst Rev* **2005**;3:CD004786.